Asymptotic behavior of the eigenvalues of a boundary value problem for the operator Schrödinger equation with a spectral parameter entering the boundary condition polynomially

Ilyas F. Hashimoglu[1]

[1]Karabuk University, Faculty of Business, Department of Business Administration, Turkey
E-mail: i.hasimoglu@karabuk.edu.tr[1]

Abstract

On the space $H_1 = L_2(H, [0,1])$, where H is a separable Hilbert space, we study the asymptotic behavior of the eigenvalues of a boundary value problem for the operator Schrödinger equation. We investigate the case when one, and the same, spectral parameter participates linearly in the equation and polynomially in the boundary condition. Asymptotic formulas are obtained for the eigenvalues of the boundary value problem under consideration.

2010 Mathematics Subject Classification. **34K08**. 34L20, 35J10, 35J25, 35P20, 35P30.
Keywords. Operator differential equations, Spectrum, Eigenvalue, Asymptotic formula, Hilbert space.

1 Introduction

In this paper, on the space $H_1 = L_2(H, [0,1])$, where H is a separable Hilbert space, we study the asymptotics of the eigenvalues of the following boundary value problem for the operator Schrödinger equation:

$$-u''(x) + \frac{\nu^2 - \frac{1}{4}}{x^2} u(x) + Au(x) = \lambda u(x), \quad x \in (0,1), \quad \nu \geq \frac{1}{2}, \tag{1.1}$$

$$u(0) = 0, \quad u'(1) + \left(\alpha_0 + \alpha_1 \lambda + \alpha_2 \lambda^2\right) u(1) = 0, \tag{1.2}$$

where λ is a spectral parameter, A is a self-adjoint, positive-definite operator in H, the inverse operator A^{-1} is completely continuous in H; $\alpha_0, \alpha_1, \alpha_2$ are real numbers: $\alpha_0 \geq 0$ and $\alpha_2 < 0$.

In [1], Hashimoglu, Akın and Mamedov investigated the discreteness and some other properties of the spectrum for the Schrödinger operator on the space $H_1 = L_2(H, [0,\infty))$.

In the papers [2] (by Gorbachuk and Rybak) and [3] (by Rybak), the asymptotic behavior of the eigenvalues of the following boundary value problem for a second-order elliptic differential operator equation

$$-u''(x) + Au(x) = \lambda u(x), \quad x \in (0,1), \tag{1.3}$$

with the boundary conditions

$$u'(0) + \lambda u(0) = 0, \quad u(1) = 0 \tag{1.4}$$

was studied on the space $H_1 = L_2(H, [0,1])$.

In [4], Aliev investigated the eigenvalues of two boundary value problems: (1.3), (1.4) and (1.3) with boundary conditions

$$u'(0) + \lambda u(0) = 0, \quad u'(1) - \lambda u(1) = 0 \tag{1.5}$$

Advanced Studies: Euro-Tbilisi Mathematical Journal 16, supplement issue 4 (2023), pp. 35–42.
DOI: 10.32513/asetmj/1932200823404
Tbilisi Centre for Mathematical Sciences.
Received by the editors: 20 January 2023.
Accepted for publication: 20 April 2023.

and proved that the set of the eigenvalues of these problems is discrete and has two series of eigenvalues: $\lambda_k \sim \sqrt{\mu_k}$ and $\lambda_{n,k} \sim n^2\pi^2 + \mu_k, \quad k, n \in N$.

In [5], similar problem for equation (1.3) with boundary conditions of the form

$$u'(0) + d\lambda^2 u(0) = 0, \quad u(1) = 0 \tag{1.6}$$

was studied.

In the paper [6], Aliev considered asymptotic behavior of eigenvalues for the equation (1.3) with the boundary conditions

$$u'(0) + \lambda^2 u(0) = 0, \quad u'(1) - \lambda^2 u(1) = 0. \tag{1.7}$$

It was shown that the problem (1.3), (1.7) has three series of eigenvalues, one of which converges to zero, and the other two asymptotically behave as $(2n)^2\pi^2 + \mu_k$ and $(2n-1)^2\pi^2 + \mu_k$, where $\mu_k = \mu_k(A)$ are the eigenvalues of the operator A, where $\mu_k = \mu_k(A)$ are the eigenvalues of the operator A.

In [7] and [8], Kapustin considered a boundary value problem for a second-order ordinary differential equation in the case when one, and the same, spectral parameter λ linearly participates in the equation and quadratically in one of boundary conditions. The asymptotic behavior of the eigenvalues of the considered boundary value problem was studied.

In [9], Mamedov proved the oscillation theorem for the problem with the spectral parameter at the boundary and obtained asymptotic formulas for the eigenvalues.

In [10, 11], Kerimov and Mamedov considered a boundary value problem for a second-order ordinary differential equation, when one, and the same, spectral parameter λ quadratically participates in the equation, while in the boundary conditions it appears as a quadratic trinomial (with respect to λ) and studied the asymptotic behavior of eigenvalues of the considered boundary value problem.

In [12], Guliyev studied various direct and inverse spectral problems for the one-dimensional Schrödinger equation with distributional potential and boundary conditions containing the eigenvalue parameter.

In [13], Aslanova considered asymptotics of the eigenvalues of boundary problem for the operator Schrödinger equation where spectral parameter participates linearly in the equation and in the boundary condition. She proved that the Schrödinger equation (1.1) with the boundary conditions

$$u(0) = 0, \quad u'(1) - hu(1) = \lambda u(1) \tag{1.8}$$

has a discrete spectrum and two series of eigenvalues:

$\lambda_k \sim -h\sqrt{\mu_k}$ and $\lambda_{m,k} \sim \left(\pi m + \frac{1}{2}\nu\pi - \frac{1}{4}\pi\right)^2 + \mu_k$.

In [14], Hashimoglu studied the asymptotic behavior of the eigenvalues of the boundary value problem for the operator Schrödinger equation (1.1) with the boundary conditions

$$u(0) = 0, \quad u'(1) - \lambda^2 u(1) = 0 \tag{1.9}$$

Note that this condition is a particular case of the condition (1.2), which we treat in the present study.

Problems such as (1.1), (1.2) and (1.1), (1.9) arise upon separation of variables in heat or wave equations, where one of the boundary conditions contains partial derivative with respect to time.

This is why it is so important to study such problems. Especially, problems with nonlinear boundary conditions are more important. The purpose of this study is to determine the asymptotics of the eigenvalues of boundary value problem for the operator Schrödinger equation in the case when one, and the same, spectral parameter participates linearly in the equation and quadratically in the boundary condition. We prove that the eigenvalues of the problem (1.1)-(1.2) are real and simple. Furthermore, it will be shown that the eigenvalues asymptotically behave as $\left(\frac{1}{2}\nu\pi + \frac{3}{4}\pi + \pi n\right)^2 + \mu_k$ or $(j_{n+1})^2 + \mu_k$, where the numbers $j_1 < j_2 < j_3 < \cdots < j_n < \cdots$ are the roots of the equation $J_\nu(z) = 0$, and $\mu_k = \mu_k(A)$ are the eigenvalues of the operator A.

2 Some properties of eigenvalues

Lemma 2.1. The eigenvalues of the boundary value problem (1.1), (1.2) are real numbers.

Proof. Let $\mu_1 \le \mu_2 \le \cdots \le \mu_n \le \cdots$ be the eigenvalues and $\{e_k\}_{k=1}^{\infty}$ be the corresponding eigenvectors of the operator A. The set $\{e_k\}_{k=1}^{\infty}$ forms a complete orthonormal basis in the space H, and for each vector-function $u(x)$ we have $u(x) = \sum_{k=1}^{\infty} (u(x), e_k)_H \cdot e_k$. For the Fourier coefficients $u_k(x) = (u(x), e_k)_H$, we get the following spectral problem

$$-u_k''(x) + \frac{\nu^2 - \frac{1}{4}}{x^2} u_k(x) + \mu_k u_k(x) = \lambda u_k(x), \quad x \in (0,1), \quad \nu \ge \frac{1}{2}, \tag{2.1}$$

$$u_k(0) = 0, \quad u_k'(1) + \left(\alpha_0 + \alpha_1\lambda + \alpha_2\lambda^2\right) u_k(1) = 0. \tag{2.2}$$

The study of the eigenvalue problem (1.1), (1.2) is reduced to the study of eigenvalues of the problem (2.1), (2.2) for different natural k. The spectrum of the problem (1.1), (1.2) consists of the union of eigenvalues of the problem (2.1), (2.2) for all natural k.

Let λ be an eigenvalue of the problem (2.1), (2.2) and $u_k(x)$ be the corresponding eigenfunction. Multiplying both sides of equality (2.1) by the function $\overline{u_k(x,\lambda)}$ and integrating the identity with respect to x from 0 to 1; we have

$$-\int_0^1 u_k''(x,\lambda)\,\overline{u_k(x,\lambda)}dx + \mu_k \int_0^1 |u_k(x,\lambda)|^2 dx + \int_0^1 \frac{\nu^2 - \frac{1}{4}}{x^2} |u_k(x,\lambda)|^2 dx$$

$$= \lambda \int_0^1 |u_k(x,\lambda)|^2 dx \tag{2.3}$$

Calculating the first integral and using boundary conditions (2.2), we get

$$-\int_0^1 u_k''(x,\lambda)\,\overline{u_k(x,\lambda)}dx = u_k'(x,\lambda)\,\overline{u_k(x,\lambda)}|_0^1 - \int_0^1 u_k'(x,\lambda)\,\overline{u_k'(x,\lambda)}dx$$

$$= u_k'(1,\lambda)\,\overline{u_k(1,\lambda)} - u_k'(0,\lambda)\,\overline{u_k(0,\lambda)} - \int_0^1 |u_k'(x,\lambda)|^2 dx$$

$$= -\left(\alpha_0 + \alpha_1\lambda + \alpha_2\lambda^2\right) |u_k(1,\lambda)|^2 - \int_0^1 |u_k'(x,\lambda).|^2 dx$$

Consequently, from (2.3), it follows that

$$-\alpha_2\lambda^2|u_k(1,\lambda)|^2+\lambda\left(\int_0^1|u_k(x,\lambda)|^2dx-\alpha_1|u_k(1,\lambda)|^2\right)-\alpha_0|u_k(1,\lambda)|^2$$

$$-\int_0^1|u_k'(x,\lambda)|^2dx-\int_0^1\frac{\nu^2-\frac{1}{4}}{x^2}|u_k(x,\lambda)|^2dx-\mu_k\int_0^1|u_k(x,\lambda)|^2dx=0. \tag{2.4}$$

Denote

$$a_k(\lambda)=-\alpha_2\ |u_k(1,\lambda)|^2,\quad b_k(\lambda)=\int_0^1|u_k(x,\lambda)|^2dx-\alpha_1|u_k(1,\lambda)|^2,$$

$$c_k(\lambda)=-\alpha_0|u_k(1,\lambda)|^2-\int_0^1|u_k'(x,\lambda)|^2dx-\int_0^1\frac{\nu^2-\frac{1}{4}}{x^2}|u_k(x,\lambda)|^2dx-\mu_k\int_0^1|u_k(x,\lambda)|^2dx. \tag{2.5}$$

From (2.4), it follows that the eigenvalue λ is a root of the equation

$$a_k(\lambda)z^2+b_k(\lambda)z+c_k(\lambda)=0,$$

for every k.
Since $a_k(\lambda)\geq 0$, $b_k(\lambda)>0$, $c_k(\lambda)<0$ for any $k\in N$, then $b_k^2(\lambda)-4a_k(\lambda)c_k(\lambda)>0$. Consequently, for every k, the problem (2.1), (2.2) has only real roots. Lemma 2.1 is proved. Q.E.D.

Lemma 2.2. The number $\lambda=0$ is not an eigenvalue of the boundary value problem (1.1), (1.2).

Proof. It is sufficient to prove that the boundary value problem (2.1), (2.2) for $\lambda=0$, i.e., the problem

$$-u_k''(x)+\frac{\nu^2-\frac{1}{4}}{x^2}u_k(x)+\mu_k u_k(x)=0,\quad x\in(0,1),\quad \nu\geq\frac{1}{2}, \tag{2.6}$$

$$u_k(0)=0,\quad u_k'(1)+\alpha_0u_k(1)=0 \tag{2.7}$$

has only trivial solution, for every natural k.

Denote by L_0 the differential operator acting in the space $L_2(0,1)$ generated by the differential expression

$$l_0[u]=-u''(x)+\frac{\nu^2-\frac{1}{4}}{x^2}u(x)+\mu_k u(x)$$

and boundary conditions

$$u(0)=0,\quad u'(1)+\alpha_0u(1)=0.$$

Consider the scalar product (L_0u, u). We have:

$$\begin{aligned}
& (L_0u, u) \\
= & -\int_0^1 u''(x)\,\overline{u(x)}dx + \int_0^1 \frac{\nu^2-\frac{1}{4}}{x^2}u(x)\,\overline{u(x)}dx + \mu_k\int_0^1 u(x)\,\overline{u(x)}dx \\
= & -u'(x)\,\overline{u(x)}|_0^1 + \int_0^1 u'(x)\,\overline{u'(x)}dx \\
& + \int_0^1 \frac{\nu^2-\frac{1}{4}}{x^2}u(x)\,\overline{u(x)}dx + \mu_k\int_0^1 u(x)\,\overline{u(x)}dx \\
= & \alpha_0|u(1)|^2 + \int_0^1 |u'(x)|^2dx + \int_0^1 \frac{\nu^2-\frac{1}{4}}{x^2}|u(x)|^2dx + \mu_k\int_0^1 |u(x)|^2dx. \qquad (2.8)
\end{aligned}$$

From (2.8), it follows that $(L_0u, u) > 0$, when $u(x) \neq 0$; and $L_0u = 0$, when $u(x) = 0$. Hence, for every natural k, the problem (2.6), (2.7) has only trivial solution. This proves the Lemma 2.2. Q.E.D.

3 Asymptotic formulas for eigenvalues

Theorem 3.1. Let A be a self-adjoint, positive-definite operator in a separable Hilbert space H and A^{-1} be completely continuous in H. Then the following asymptotic expression holds for the eigenvalues of the boundary value problem (1.1), (1.2):

$$\lambda_{k,n} \sim \mu_k + \left(\frac{1}{2}\nu\pi + \frac{3}{4}\pi + \pi n\right)^2 = \mu_k + (j_{n+1})^2,$$

where the numbers $j_1 < j_2 < j_3 < \cdots < j_n < \cdots$ are the roots of the equation $J_\nu(z) = 0$, and $\mu_k = \mu_k(A) \to \infty$ (as $k \to \infty$) are the eigenvalues of the operator A.

Proof. The general solution of the differential equations (2.1) is of the form

$$u_k(x,\lambda) = k_1\sqrt{2x}\cdot J_\nu\left(\sqrt{\lambda-\mu_k}x\right) + k_2\sqrt{2x}\cdot Y_\nu\left(\sqrt{\lambda-\mu_k}x\right), \qquad (3.1)$$

where k_1 and k_2 are arbitrary constants; $J_\nu(z)$ and $Y_\nu(z)$ are the Bessel functions of first and second kinds, respectively. From the first boundary condition $u_k(0,\lambda) = 0$ it follows that $k_2 = 0$. Therefore,

$u_k(x,\lambda) = k_1\sqrt{2x}\cdot J_\nu\left(\sqrt{\lambda-\mu_k}x\right)$.

From the second condition of (2.2) we get

$$u_k'(1,\lambda) + \left(\alpha_0 + \alpha_1\lambda + \alpha_2\lambda^2\right)u_k(1,\lambda) =$$

$$\left(\frac{1}{\sqrt{2}} - \sqrt{2}\nu\right)J_\nu\left(\sqrt{\lambda-\mu_k}\right) + \sqrt{2(\lambda-\mu_k)}J_{\nu-1}\left(\sqrt{\lambda-\mu_k}\right) - \sqrt{2}\lambda^2 J_\nu\left(\sqrt{\lambda-\mu_k}\right) = 0,$$

or

$$\left(1 - 2\nu + 2\alpha_0 + 2\alpha_1\lambda + 2\alpha_2\lambda^2\right)\cdot J_\nu\left(\sqrt{\lambda-\mu_k}\right) + 2\sqrt{\lambda-\mu_k}\cdot J_{\nu-1}\left(\sqrt{\lambda-\mu_k}\right) = 0. \qquad (3.2)$$

Denote $\sqrt{\lambda-\mu_k}=y$. Hence, $\lambda=\mu_k+y^2$, $\lambda^2=\left(\mu_k+y^2\right)^2$ and $y\to\infty$ as $\lambda\to\infty$. Then equation (3.2) takes the form

$$\left(1-2\nu+2\alpha_1\left(\mu_k+y^2\right)+2\alpha_2\left(\mu_k+y^2\right)^2\right)\cdot J_\nu(y)+2y\cdot J_{\nu-1}(y)=0. \tag{3.3}$$

Using the asymptotic expression for $J_\nu(y)$ at $y\to\infty$ we obtain

$$\begin{aligned}&\left(1-2\nu+2\alpha_1\left(\mu_k+y^2\right)+2\alpha_2\left(\mu_k+y^2\right)^2\right)\cdot\\&\left[\cos\left(y-\frac{1}{2}\nu\pi-\frac{1}{4}\pi\right)\left(1-\frac{c_1}{y^2}+\frac{c_2}{y^4}+O\left(\frac{1}{y^6}\right)\right)\right.\\&\left.-\sin\left(y-\frac{1}{2}\nu\pi-\frac{1}{4}\pi\right)\left(\frac{4\nu^2-1}{8y}-\frac{c_3}{y^3}+O\left(\frac{1}{y^5}\right)\right)\right]\\&-2y\left[\sin\left(y-\frac{1}{2}\nu\pi-\frac{1}{4}\pi\right)\left(1+O\left(\frac{1}{y^2}\right)\right)\right.\\&\left.+\cos\left(y-\frac{1}{2}\nu\pi-\frac{1}{4}\pi\right)\left(\frac{4\nu^2-1}{8y}+O\left(\frac{1}{y^3}\right)\right)\right]=0,\end{aligned} \tag{3.4}$$

where c_1, c_2 and c_3 are some positive real constants.

Denote $y-\frac{1}{2}\nu\pi-\frac{1}{4}\pi=z$. After some processing we get

$$\begin{aligned}&\cos(z)\left[\left(1-2\nu+2\alpha_1\left(\mu_k+y^2\right)+2\alpha_2\left(\mu_k+y^2\right)^2\right)\left(1-\frac{c_1}{y^2}+\frac{c_2}{y^4}+O\left(\frac{1}{y^6}\right)\right)+\frac{4\nu^2-1}{4}+O\left(\frac{1}{y^2}\right)\right]\\&-\sin(z)\left[\left(1-2\nu+2\alpha_1\left(\mu_k+y^2\right)+2\alpha_2\left(\mu_k+y^2\right)^2\right)\left(\frac{4\nu^2-1}{8y}-\frac{c_3}{y^3}+O\left(\frac{1}{y^5}\right)\right)+2y+O\left(\frac{1}{y}\right)\right]=0.\end{aligned} \tag{3.5}$$

Equation (3.5) takes the form

$$\tan(z)=\frac{\left(1-2\nu+2\alpha_1\left(\mu_k+y^2\right)+2\alpha_2\left(\mu_k+y^2\right)^2\right)\cdot\left(1-\frac{c_1}{y^2}+\frac{c_2}{y^4}+O\left(\frac{1}{y^6}\right)\right)+\frac{4\nu^2-1}{4}+O\left(\frac{1}{y^2}\right)}{\left(1-2\nu+2\alpha_1\left(\mu_k+y^2\right)+2\alpha_2\left(\mu_k+y^2\right)^2\right)\cdot\left(\frac{4\nu^2-1}{8y}-\frac{c_3}{y^3}+O\left(\frac{1}{y^5}\right)\right)+2y+O\left(\frac{1}{y}\right)}$$

When $y\to\infty$,

$$\tan(z)=\frac{\left(1-2\nu+2\alpha_1\left(\mu_k+y^2\right)+2\alpha_2\left(\mu_k+y^2\right)^2\right)\left(1+O\left(\frac{1}{y^2}\right)\right)}{\left(1-2\nu+2\alpha_1\left(\mu_k+y^2\right)+2\alpha_2\left(\mu_k+y^2\right)^2\right)\left(\frac{4\nu^2-1}{8y}+O\left(\frac{1}{y^3}\right)\right)}=\frac{8y}{4\nu^2-1}+O\left(\frac{1}{y^2}\right)$$

Equation $\tan(z)=\frac{8y}{4\nu^2-1}+O\left(\frac{1}{y^2}\right)$ has roots $z_k=y_k-\frac{1}{2}\nu\pi-\frac{1}{4}\pi\sim\frac{\pi}{2}+\pi k$, $k\in Z^+$ when $y\to\infty$. From here: $y_k=\sqrt{\lambda-\mu_k}=\frac{1}{2}\nu\pi+\frac{3}{4}\pi+\pi k$ or

$$\lambda_{k,n}\sim\mu_k+\left(\frac{1}{2}\nu\pi+\frac{3}{4}\pi+\pi n\right)^2. \tag{3.6}$$

Let numbers $j_1 < j_2 < j_3 < \cdots < j_n < \cdots$ satisfy the equality $J_\nu(j_n) = 0$ for $n = 1, 2, 3, \ldots$.

Taking into account the asymptotics of numbers j_n, i.e. $j_n \sim \frac{1}{2}\nu\pi - \frac{1}{4}\pi + \pi n$, for $n \to \infty$, from (3.6) we obtain the following evaluation:

$$\lambda_{k,n} \sim \mu_k + (j_{n+1})^2. \tag{3.7}$$

Theorem 3.1 is proved. Q.E.D.

References

[1] I. Hashimoglu, Ö. Akın and K.R. Mamedov, *The discreteness of the spectrum of the Schrödinger operator equation and some properties of the s-numbers of the inverse Schrödinger operator,* Mathematical Methods in the Applied Science, Vol. 42, No. 7 (2019), pp. 2231–2243

[2] V.I. Gorbachuk and M.A. Rybak, *On the boundary-value problems for the Sturm-Liouville operator equation with a spectral parameter in the equation and in the boundary condition,* Direct and Inverse Problems of the Scattering Theory, Kiev (1981), pp. 3–16 (Russian)

[3] M.A. Rybak, *Asymptotic distribution of the eigenvalues of some boundary value problems for Sturm-Liouville operator equations,* Ukrainian Mathematical Journal, Vol. 32, No. 2 (1980), pp. 159–162

[4] B.A. Aliev, *Asymptotic behavior of the eigenvalues of a boundary value problem for a second-order elliptic differential operator equation,* Ukrainian Mathematical Journal, Vol. 58, No. 8 (2006), pp. 1298–1306

[5] B.A. Aliev, *Asymptotic behavior of eigenvalues of a boundary value problem for a second-order elliptic differential-operator equation with spectral parameter quadratically occurring in the boundary condition*, Differential Equations, Vol. 54, No. 9 (2018), pp. 1256–1260

[6] B.A. Aliev, *On eigenvalues of a boundary value problem for a second order elliptic differential-operator equation,* Proceedings of the Institute of Mathematics and Mechanics, National Academy of Sciences of Azerbaijan, Vol. 45, No. 2 (2019), pp. 213–221

[7] N.Yu. Kapustin, *On a spectral problem in the theory of the heat operator,* Differential Equations, Vol. 45, No. 10 (2009), pp. 1509–1511

[8] N.Yu. Kapustin, *On the uniform convergence in C1 of Fourier series for a spectral problem with squared spectral parameter in a boundary condition,* Differential Equations, Vol. 47, No. 10 (2011), pp. 1408–1413

[9] Kh.R. Mamedov, *On boundary value problem with parameter in boundary conditions,* Spectral Theory of Operators and Its Applications, Vol. 11 (1997), pp. 117–121

[10] N.B. Kerimov and Kh.R. Mamedov, *On a boundary value problem with a spectral parameter in the boundary conditions,* Siberian Mathematical Journal, Vol. 40, No. 2 (1999), pp. 281–290

[11] N.B. Kerimov and Kh.R. Mamedov, *The Sturm-Liouville problem with non-linear spectral parameter in the boundary conditions,* Transactions of National Academy of Sciences of Azerbaijan, Series of Physical-Technical and Mathematical Sciences, Issue Mathematics, Vol. 21, No. 4 (2001), pp. 100–104

[12] N.J. Guliyev, *Schrödinger operators with distributional potentials and boundary conditions dependent on the eigenvalue parameter,* Journal of Mathematical Physics, Vol. 60 (2019), Article 063501. DOI: 10.1063/1.5048692

[13] N.M. Aslanova, *Study of the asymptotic eigenvalue distribution and trace formula of a second order operator-differential equation,* Boundary Value Problems, Vol. 2011 (2011), Article 7, pp. 1–22. DOI: 10.1186/1687-2770-2011-7

[14] I.F. Hashimoglu, *Asymptotics of the eigenvalues of a boundary value problem for the operator Schrödinger equation with boundary conditions nonlinearly dependent on the spectral parameter,* Journal of Samara State Technical University, Ser. Physical and Mathematical Sciences, Vol. 25, No. 4 (2021), pp. 607–615. DOI: 10.14498/vsgtu1894