Certain subclass of meromorphic functions involving generalized differential operator

B. Venkateswarlu[1], P. Thirupathi Reddy[2] and N. Rani[3]

[1]Department of Mathematics, GSS, GITAM University, Doddaballapur-562 163, Bengaluru Rural, Karnataka, India.
[2]Department of Mathematics, Kakatiya Univeristy, Warangal- 506 009, Telangana, India.
[3]Department of Mathematics, PRIME College , Modavalasa - 534 002, Visakhapatnam, A. P., India.

E-mail: bvlmaths@gmail.com[1], reddypt2@gmail.com[2], raninekkanti1111@gmail.com[3]

Abstract

In this paper we introduce and study a new subclass of meromorphically uniformly convex functions with positive coefficients defined by a differential operator and obtain coefficient estimates, growth and distortion theorem, radius of convexity,integral transforms, convex linear combinations, convolution properties and δ-neighborhoods for the class $\Sigma_p(\alpha, \beta)$.

2010 Mathematics Subject Classification. **30C45**
Keywords. uniformly convex, uniformly starlike, coefficient estimates.

1 Introduction

Let A denote the class of all functions $f(z)$ of the form

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n \tag{1.1}$$

in the open unit disc $E = \{z \in \mathcal{C} : |z| < 1\}$. Let S be the subclass of A consisting of univalent functions and satisfy the following usual normalization condition $f(0) = f'(0) - 1 = 0$. We denote by S the subclass of A consisting of functions $f(z)$ which are all univalent in E. A function $f \in A$ is a starlike function by the order $\alpha,\ 0 \leq \alpha < 1$, if it satisfy

$$Re\left\{\frac{zf'(z)}{f(z)}\right\} > \alpha \ (z \in E). \tag{1.2}$$

We denote this class with $S^*(\alpha)$.

A function $f \in A$ is a convex function by the order $\alpha,\ 0 \leq \alpha < 1$, if it satisfy

$$Re\left\{1 + \frac{zf''(z)}{f'(z)}\right\} > \alpha \ (z \in E). \tag{1.3}$$

We denote this class with $K(\alpha)$.

Let T denote the class of functions analytic in E that are of the form

$$f(z) = z - \sum_{n=2}^{\infty} a_n z^n \quad (a_n \geq 0 \ z \in E) \tag{1.4}$$

Tbilisi Mathematical Journal 13(4) (2020), pp. 1–11.
Tbilisi Centre for Mathematical Sciences.
Received by the editors: 13 January 2019.
Accepted for publication: 10 September 2020 .

and let $T^*(\alpha) = T \cap S^*(\alpha),\ C(\alpha) = T \cap K(\alpha)$. The class $T^*(\alpha)$ and allied classes possess some interesting properties and have been extensively studied by Silverman [19] and others.

A function $f \in A$ is said to be in the class of uniformly convex functions of order γ and type β, denoted by $UCV(\beta, \gamma)$, if

$$Re\left\{1 + \frac{zf''(z)}{f'(z)} - \gamma\right\} > \beta\left|\frac{zf''(z)}{f'(z)}\right|, \tag{1.5}$$

where $\beta \geq 0, \gamma \in [-1, 1)$ and $\beta + \gamma \geq 0$ and it is said to be in the class corresponding class denoted by $SP(\beta, \gamma)$, if

$$Re\left\{\frac{zf'(z)}{f(z)} - \gamma\right\} > \beta\left|\frac{zf'(z)}{f(z)} - 1\right|, \tag{1.6}$$

where $\beta \geq 0, \gamma \in [-1, 1)$ and $\beta + \gamma \geq 0$. Indeed it follows from (1.5) and (1.6) that

$$f \in UCV(\gamma, \beta) \Leftrightarrow zf' \in SP(\gamma, \beta). \tag{1.7}$$

For $\beta = 0$, we get respectively, the classes $K(\gamma)$ and $S^*(\gamma)$. The function of the class $UCV(1, 0) \equiv UCV$ are called uniformly convex functions were introduced and studied by Goodman with geometric interpretation in [5, 6]. The class $SP(1, 0) \equiv SP$ is defined by Rønning [15]. The classes $UCV(1, \gamma) \equiv UCV(\gamma)$ and $SP(1, \gamma) \equiv SP(\gamma)$ are investigated by Rønning in [14]. For $\gamma = 0$, the classes $UCV(\beta, 0) \equiv \beta - UCV$ and $SP(\beta, 0) \equiv \beta - SP$ are defined respectively, by Kanas and Wisniowska in [8, 9].

Further Ahuja et al. [1], Bharathi et al. [2], Murugusundaramoorthy and Magesh [12] and others [10, 11, 20] have studied and investigated interesting properties for the classes $UCV(\beta, \gamma)$ and $SP(\beta, \gamma)$.

Let Σ denote the class of functions of the form be

$$f(z) = \frac{1}{z} + \sum_{m=1}^{\infty} a_m z^m \tag{1.8}$$

which are regular in domain $E = \{z : 0 < |z| < 1\}$ with a simple pole at the origin with residue 1 there. Let $\Sigma_s, \Sigma^*(\alpha)$ and $\Sigma_k(\alpha)$ $(0 \leq \alpha < 1)$ denote the subclasses of Σ that are univalent, meromorphically starlike of order α and meromorphically convex of order α respectively. Analytically $f(z)$ of the form (1.8) is in $\Sigma^*(\alpha)$ if and only if

$$Re\left\{-\frac{zf'(z)}{f(z)}\right\} > \alpha \quad (z \in E). \tag{1.9}$$

Similarly, $f \in \Sigma_k(\alpha)$ if and only if, $f(z)$ is of the form (1.8) and satisfies

$$Re\left\{-\left(1 + \frac{zf''(z)}{f'(z)}\right)\right\} > \alpha \quad (z \in E). \tag{1.10}$$

It being understood that if $\alpha = 1$ then $f(z) = \frac{1}{z}$ is the only function which is $\Sigma^*(1)$ and $\Sigma_k(1)$. The classes $\Sigma^*(\alpha)$ and $\Sigma_k(\alpha)$ have been extensively studied by Pommerenke [13], Clunie [3], Royster [17] and others.

Since, to a certain extent the work in the meromorphic univalent case has paralleled that of regular univalent case, it is natural to search for a subclass of Σ_s that has properties analogous to those of $T^*(\alpha)$. Juneja and Reddy [7] introduced the class Σ_p of functions of the form

$$f(z) = \frac{1}{z} + \sum_{m=1}^{\infty} a_m z^m \ (a_m \geq 0) \tag{1.11}$$

$$\Sigma_p^*(\alpha) = \Sigma_p \cap \Sigma^*(\alpha).$$

For functions $f(z)$ in the class Σ_p, we define a linear operator D^n by the following form

$$\begin{aligned} D^0 f(z) &= f(z) \\ D^1 f(z) &= \frac{1}{z} + 3a_1 z + 4a_2 z^2 + \cdots \ = \frac{(z^2 f(z))'}{z} \\ D^2 f(z) &= D(D^1 f(z)) \\ &\vdots \end{aligned}$$

$$D^n f(z) = D(D^{n-1} f(z)) = \frac{1}{z} + \sum_{m=1}^{\infty} (m+2)^n a_m z^m = \frac{(z^2 D^{n-1} f(z))'}{z}, \text{ for } n = 1, 2, \cdots \tag{1.12}$$

Now, we define a new subclass $\Sigma_p(\alpha, \beta)$ of Σ_p.

Definition 1.1. For $-1 \leq \alpha < 1$ and $\beta \geq 1$, we let $\Sigma_p(\alpha, \beta)$ be the subclass of Σ_p consisting of the form (1.11) and satisfying the analytic criterion

$$-Re\left\{\frac{z(D^n f(z))'}{D^n f(z)} + \alpha\right\} > \beta \left|\frac{z(D^n f(z))'}{D^n f(z)} + 1\right|, \tag{1.13}$$

$D^n f(z)$ is given by (1.12)

The main object of the paper is to study some usual properties of the geometric function theory such as coefficient bounds, growth and distortion properties, radius of convexity, convex linear combination and convolution properties, integral operators and $\delta-$neighbourhoods for the class$\Sigma_p(\alpha, \beta)$.

2 Coefficient inequality

In this section we obtain the coefficient bounds of function $f(z)$ for the class $\Sigma_p(\alpha, \beta)$.

Theorem 2.1. *A function $f(z)$ of the form (1.11) is in $\Sigma_p(\alpha, \beta)$ if*

$$\sum_{m=1}^{\infty} (m+2)^n [(1+\beta)(m+1) + 1 - \alpha] \ |a_m| \ \leq \ (1-\alpha) \ (-1 \leq \alpha < 1 \ and \ \beta \geq 1). \tag{2.1}$$

Proof. It sufficient to show that

$$\beta\left|\frac{z(D^nf(z))'}{D^nf(z)}+1\right|+Re\left\{\frac{z(D^nf(z))'}{D^nf(z)}+1\right\}\leq(1-\alpha).$$

$$\begin{aligned}\text{We have }&\beta\left|\frac{z(D^nf(z))'}{D^nf(z)}+1\right|+Re\left\{\frac{z(D^nf(z))'}{D^nf(z)}+1\right\}\\&\leq(1+\beta)\left|\frac{z(D^nf(z))'}{D^nf(z)}+1\right|\\&\leq\frac{(1+\beta)\sum\limits_{m=1}^{\infty}(m+2)^n(m+1)|a_m||z^m|}{\frac{1}{|z|}-\sum\limits_{m=1}^{\infty}(m+2)^n|a_m||z^m|}\end{aligned}$$

$$\text{Letting } z\to 1 \text{ along the real axis, we obtain}$$

$$\leq\frac{(1+\beta)\sum\limits_{m=1}^{\infty}(m+2)^n(m+1)|a_m|}{1-\sum\limits_{m=1}^{\infty}(m+2)^n|a_m|}.$$

The above expression is bounded by $(1-\alpha)$ if

$$\sum_{m=1}^{\infty}(m+2)^n[(1+\beta)(m+1)+1-\alpha]\ |a_m|\ \leq\ (1-\alpha).$$

Hence the theorem is completed. Q.E.D.

Corollary 2.2. *Let the function $f(z)$ defined by (1.11) be in the class $\Sigma_p(\alpha,\beta)$. Then*

$$a_m\leq\frac{(1-\alpha)}{\sum\limits_{m=1}^{\infty}(m+2)^n[(1+\beta)(m+1)+1-\alpha]}\quad(m\geq1).\tag{2.2}$$

Equality holds for the function of the form

$$f_m(z)=\frac{1}{z}+\frac{(1-\alpha)}{(m+2)^n[(1+\beta)(m+1)+1-\alpha]}\ z^m.\tag{2.3}$$

3 Distortion theorems

In this section we obtain Distortion bounds for the class $\Sigma_p(\alpha,\beta)$.

Theorem 3.1. *Let the function $f(z)$ defined by (1.11) be in the class $\Sigma_p(\alpha,\beta)$. Then for $0<|z|=r<1$,*

$$\frac{1}{r}-\frac{(1-\alpha)}{3^n[3+2\beta-\alpha]}\ r\leq|f(z)|\leq\frac{1}{r}+\frac{(1-\alpha)}{3^n[3+2\beta-\alpha]}r\tag{3.1}$$

with equality for the function

$$f(z)=\frac{1}{z}+\frac{(1-\alpha)}{3^n[3+2\beta-\alpha]}\ z\ \ at\ \ z=r,ir.\tag{3.2}$$

Proof. Suppose $f(z)$ is in $\Sigma_p(\alpha,\beta)$. In view of Theorem 2.1, we have
$3^n[3+2\beta-\alpha]\sum\limits_{m=1}^{\infty} a_m \leq \sum\limits_{m=1}^{\infty}(m+2)^n[(1+\beta)(m+1)+1-\alpha] \leq (1-\alpha)$
which evidently yields $\sum\limits_{m=1}^{\infty} a_m \leq \frac{1-\alpha}{3^n[3+2\beta-\alpha]}$.
Consequently, we obtain

$$\begin{aligned}|f(z)| &= \left|\frac{1}{z}+\sum_{m=1}^{\infty} a_m z^m\right| \leq \left|\frac{1}{z}\right| + \sum_{m=1}^{\infty} a_m |z|^m \\ &\leq \frac{1}{r} + r\sum_{m=1}^{\infty} a_m \\ &\leq \frac{1}{r} + \frac{1-\alpha}{3^n[3+2\beta-\alpha]}\ r.\end{aligned}$$

$$\begin{aligned}\text{Also, } |f(z)| &= \left|\frac{1}{z}+\sum_{m=1}^{\infty} a_m z^m\right| \geq \left|\frac{1}{z}\right| - \sum_{m=1}^{\infty} a_m |z|^m \\ &\geq \frac{1}{r} - r\sum_{m=1}^{\infty} a_m \\ &\geq \frac{1}{r} - \frac{1-\alpha}{3^n[3+2\beta-\alpha]}\ r.\end{aligned}$$

Hence the results (3.1) follow. Q.E.D.

Theorem 3.2. *Let the function $f(z)$ defined by* (1.11) *be in the class $\Sigma_p(\alpha,\beta)$. Then for $0<|z|=r<1$,*

$$\frac{1}{r^2} - \frac{1-\alpha}{3^n[3+2\beta-\alpha]} \leq |f'(z)| \leq \frac{1}{r^2} + \frac{1-\alpha}{3^n[3+2\beta-\alpha]}.$$

The result is sharp, the extremal function being of the form (2.3)

Proof. From Theorem 2.1, we have
$3^n[3+2\beta-\alpha]\sum\limits_{m=1}^{\infty} ma_m \leq \sum\limits_{m=1}^{\infty}(m+2)^n[(1+\beta)(m+1)+1-\alpha] \leq (1-\alpha)$ which evidently yields
$\sum\limits_{m=1}^{\infty} ma_m \leq \frac{1-\alpha}{3^n[3+2\beta-\alpha]}$.

Consequently, we obtain

$$\begin{aligned} |f'(z)| &\leq \left|\frac{1}{r^2}+\sum_{m=1}^{\infty} m a_m r^{m-1}\right| \\ &\leq \frac{1}{r^2}+\sum_{m=1}^{\infty} m a_m \\ &\leq \frac{1}{r^2}+\frac{(1-\alpha)}{3^n[3+2\beta-\alpha]}. \\ \text{Also, } |f'(z)| &\geq \left|\frac{1}{r^2}-\sum_{m=1}^{\infty} m a_m r^{m-1}\right| \\ &\geq \frac{1}{r^2}\ -\sum_{m=1}^{\infty} m a_m \\ &\geq \frac{1}{r^2}+\frac{(1-\alpha)}{(3^n[3+2\beta-\alpha]}. \end{aligned}$$

This completes the proof. Q.E.D.

4 Class preserving integral operators

In this section we consider the class preserving integral operator of the form (1.11).

Theorem 4.1. *Let the function $f(z)$ defined by (1.11) be in the class $\Sigma_p(\alpha,\beta)$. Then*

$$f(z)=cz^{-c-1}\int_0^z t^c f(t)dt=\frac{1}{z}+\sum_{m=1}^{\infty}\frac{c}{c+m+1}\ a_m z^m \quad (c>0) \tag{4.1}$$

is in $\Sigma_p(\delta,\beta)$, where

$$\delta(\alpha,\beta,c)=\frac{(m+1)(1+\beta)+\alpha c(1-\beta)+(1-\alpha)}{.}(c+m+1)(1+\beta)+(1-\alpha) \tag{4.2}$$

The result is sharp for $f(z)=\frac{1}{z}+\frac{(1-\alpha)}{3^n[3+2\beta-\alpha]}z$.

Proof. Suppose $f(z)=\frac{1}{z}+\sum\limits_{m=1}^{\infty} a_m z^m$ is in $\Sigma_p(\alpha,\beta)$. We have

$f(z)=cz^{-c-1}\int\limits_0^z t^c f(t)dt=\frac{1}{z}+\sum\limits_{m=1}^{\infty}\frac{c}{c+m+1}\ a_m z^m \quad (c>0)$.

It is sufficient to show that

$$\sum_{m=1}^{\infty}\frac{(m+2)^n[(1+\beta)(m+1)+1-\delta]}{1-\delta}\frac{c}{c+m+1}a_m\leq 1. \tag{4.3}$$

Since $f(z)$ is in $\Sigma_p(\alpha,\beta)$, we have

$$\sum_{m=1}^{\infty}\frac{(m+2)^n[(1+\beta)(m+1)+1-\alpha]}{1-\alpha}|a_m|\leq 1. \tag{4.4}$$

Thus (4.3) will be satisfied if
$\sum_{m=1}^{\infty} \frac{[(1+\beta)(m+1)+1-\delta]}{1-\delta} \frac{c}{c+m+1} \leq \sum_{m=1}^{\infty} \frac{[(1+\beta)(m+1)+1-\alpha]}{1-\alpha}$.
Solving for δ, we obtain

$$\delta \leq \frac{(1+\beta)(m+1)+\alpha c(1-\beta)+(1-\alpha)}{(c+m+1)(1-\beta)+(1-\alpha)} = G(m). \tag{4.5}$$

A simple computation will show that $G(m)$ is increasing and $G(m) \geq G(1)$. Using this, the result follows. Q.E.D.

5 Convex linear combinations and convolution properties

In this section we obtain sharp for $f(z)$ is meromorphically convex of order δ and necessary and sufficient condition for $f(z)$ is in the class $\Sigma_p(\alpha,\beta)$ and also proved that convolution is in the class.

Theorem 5.1. *If the function* $f(z) = \frac{1}{z} + \sum_{m=1}^{\infty} a_m z^m$ *is in* $\Sigma_p(\alpha,\beta)$ *then* $f(z)$ *is meromorphically convex of order* $\delta(0 \leq \delta < 1)$ *in* $|z| < r = r(\alpha,\beta,\delta)$ *where*

$$r(\alpha,\beta,\delta) = \inf_{n\geq 1} \left\{ \frac{(1-\delta)(m+2)^n[(1+\beta)(1+m)+1-\alpha]}{(1-\alpha)m(m+2-\delta)} \right\}^{\frac{1}{m+1}}.$$

The result is sharp.

Proof. Let $f(z)$ be in $\Sigma_p(\alpha,\beta)$. Then, by Theorem 2.1, we have

$$\sum_{m=1}^{\infty} (m+2)^n[(1+\beta)(1+m)+1-\alpha]|a_m| \leq (1-\alpha). \tag{5.1}$$

It is sufficient to show that $\left|2+\frac{zf''(z)}{f'(z)}\right| \leq (1-\delta)$ for $|z| < r = r(\alpha,\beta,\delta)$, where $r(\alpha,\beta,\delta)$ is specified in the statement of the theorem. Then

$$\left|2+\frac{zf''(z)}{f'(z)}\right| = \left|\frac{\sum_{m=1}^{\infty} m(m+1)a_m z^{m-1}}{\frac{-1}{z^2}+\sum_{m=1}^{\infty} m a_m z^{m-1}}\right| \leq \frac{\sum_{m=1}^{\infty} m(m+1)a_m|z|^{m+1}}{1-\sum_{m=1}^{\infty} m a_m|z|^{m+1}}.$$

This will be bounded by $(1-\delta)$ if

$$\sum_{m=1}^{\infty} \frac{m(m+2-\delta)}{1-\delta} a_m|z|^{m+1} \leq 1. \tag{5.2}$$

By (5.1), it follows that (5.2) is true if

$$\frac{m(m+2-\delta)}{1-\delta}|z|^{m+1} \leq \frac{(m+2)^n[(1+\beta)(m+1)+1-\alpha]}{1-\alpha}|a_m| \quad (m \geq 1)$$

$$\text{or} \quad |z| \leq \left\{ \frac{(1-\delta)(m+2)^n[(1+\beta)(m+1)+1-\alpha]}{(1-\alpha)m(m+2-\delta)} \right\}^{\frac{1}{m+1}}. \tag{5.3}$$

Setting $|z| = r(\alpha, \beta, \delta)$ in (5.3), the result follows. The result is sharp for the function

$$f_m(z) = \frac{1}{z} + \frac{(1-\alpha)}{(m+2)^n[(1+\beta)(m+1)+1-\alpha]} z^m \quad (m \geq 1).$$

Q.E.D.

Theorem 5.2. *Let* $f_0(z) = \frac{1}{z}$ *and* $f_m(z) = \frac{1}{z} + \frac{(1-\alpha)}{(m+2)^n[(1+\beta)(m+1)+1-\alpha]} z^m \quad (m \geq 1)$. *Then* $f(z) = \frac{1}{z} + \sum_{m=1}^{\infty} a_m z^m$ *is in the class* $\Sigma_p(\alpha, \beta)$ *if and only if it can be expressed in the form* $f(z) = \omega_0 f_0(z) + \sum_{m=1}^{\infty} \omega_m f_m(z)$, *where* $\omega_0 \geq 0, \omega_m \geq 0, m \geq 1$ *and* $\omega_0 + \sum_{m=1}^{\infty} \omega_m = 1$.

Proof. Let $f(z) = \omega_0 f_0(z) + \sum_{m=1}^{\infty} \omega_m f_m(z)$ with $\omega_0 \geq 0, \omega_m \geq 0, m \geq 1$ and $\omega_0 + \sum_{m=1}^{\infty} \omega_m = 1$. Then

$$\begin{aligned} f(z) &= \omega_0 f_0(z) + \sum_{m=1}^{\infty} \omega_m f_m(z) \\ &= \frac{1}{z} + \sum_{m=1}^{\infty} \omega_m \frac{(1-\alpha)}{(m+2)^n[(1+\beta)(m+1)+1-\alpha]} z^m \\ \text{Since} \sum_{m=1}^{\infty} &\frac{[(m+2)^n[(1+\beta)(m+1)+1-\alpha]}{(1-\alpha)} \omega_m \frac{(1-\alpha)}{(m+2)^n[(1+\beta)(m+1)+1-\alpha]} \\ &= \sum_{m=1}^{\infty} \omega_m = 1 - \omega_0 \leq 1. \end{aligned}$$

By Theorem 2.1, $f(z)$ is in the class $\Sigma_p(\alpha, \beta)$.

Conversely suppose that the function $f(z)$ is in the class $\Sigma_p(\alpha, \beta)$, since

$$a_m \leq \frac{(1-\alpha)}{(m+2)^n[(1+\beta)(m+1)+1-\alpha]} z^m \quad (m \geq 1).$$

$$\omega_m = \sum_{m=1}^{\infty} \frac{(m+2)^n[(1+\beta)(m+1)+1-\alpha]}{(1-\alpha)} a_m \quad \text{and} \quad \omega_0 = 1 - \sum_{m=1}^{\infty} \omega_m.$$

It follows that $f(z) = \omega_0 f_0(z) + \sum_{m=1}^{\infty} \omega_m f_m(z)$.

This completes the proof of the theorem. Q.E.D.

For the functions $f(z) = \frac{1}{z} + \sum_{m=1}^{\infty} a_m z^m$ and $g(z) = \frac{1}{z} + \sum_{m=1}^{\infty} b_m z^m$ belongs to Σ_p, we denoted by $(f * g)(z)$ the convolution of $f(z)$ and $g(z)$ and defined as

$$(f * g)(z) = \frac{1}{z} + \sum_{m=1}^{\infty} a_m b_m z^m.$$

Theorem 5.3. *If the function* $f(z) = \frac{1}{z} + \sum_{m=1}^{\infty} a_m z^m$ *and* $g(z) = \frac{1}{z} + \sum_{m=1}^{\infty} b_m z^m$ *are in the class* $\Sigma_p(\alpha, \beta)$ *then* $(f * g)(z)$ *is in the class* $\Sigma_p(\alpha, \beta)$.

Proof. Suppose $f(z)$ and $g(z)$ are in $\Sigma_p(\alpha, \beta)$. By Theorem 2.1, we have

$$\sum_{m=1}^{\infty} \frac{(m+2)^n[(1+\beta)(m+1)+1-\alpha]}{(1-\alpha)} a_m \leq 1$$

$$\text{and} \sum_{m=1}^{\infty} \frac{(m+2)^n[(1+\beta)(m+1)+1-\alpha]}{(1-\alpha)} b_m \leq 1 \ .$$

Since $f(z)$ and $g(z)$ are regular are in E, so is $(f * g)(z)$. Further more

$$\begin{aligned}
&\sum_{m=1}^{\infty} \frac{(m+2)^n[(1+\beta)(m+1)+1-\alpha]}{(1-\alpha)} a_m b_m \\
&\leq \sum_{m=1}^{\infty} \left\{ \frac{(m+2)^n[(1+\beta)(m+1)+1-\alpha]}{(1-\alpha)} \right\}^2 a_m b_m \\
&\leq \left(\sum_{m=1}^{\infty} \frac{(m+2)^n[(1+\beta)(m+1)+1-\alpha]}{(1-\alpha)} a_m \right) \\
&\qquad \left(\sum_{m=1}^{\infty} \frac{(m+2)^n[(1+\beta)(m+1)+1-\alpha]}{(1-\alpha)} b_m \right) \\
&\leq 1.
\end{aligned}$$

Hence, by Theorem 2.1, $(f * g)(z)$ is in the class $\Sigma_p(\alpha, \beta)$. Q.E.D.

6 Neighborhoods for the class $\Sigma_p(\alpha, \beta, \gamma)$

In this section we define the $\delta-$neighborhood of a function $f(z)$ and establish a relation between $\delta-$neighborhood and $\Sigma_p((\alpha, \beta, \gamma)$ class of a function.

Definition 6.1. A function $f \in \Sigma_p$ is said to in the class $\Sigma_p(\alpha, \beta, \gamma)$ if there exists a function $g \in \Sigma_p(\alpha, \beta)$ such that

$$\left| \frac{f(z)}{g(z)} - 1 \right| < (1-\gamma) \quad (z \in E,\ 0 \leq \gamma < 1). \tag{6.1}$$

Following the earlier works on neighborhoods of analytic functions by Goodman [4] and Ruschweyh [18]. We defined the $\delta-$neighborhood of a function $f \in \Sigma_p$ by

$$N_\delta(f) = \left\{ g \in \Sigma_p \ \mid \ g(z) = \frac{1}{z} + \sum_{m=1}^{\infty} b_m z^m \ : \ \sum_{m=1}^{\infty} m|a_m - b_m| \leq \delta \right\}. \tag{6.2}$$

Theorem 6.2. *If* $g \in \Sigma_p(\alpha, \beta)$ *and*

$$\gamma = 1 - \frac{\delta[3 + 2\beta - \alpha]}{2(1+\beta)} \tag{6.3}$$

then $N_\delta(g) \subset \Sigma_p(\alpha, \beta, \gamma)$.

Proof. Let $f \in N_\delta(g)$. Then we find from (6.2) that

$$\sum_{m=1}^{\infty} m|a_m - b_m| \leq \delta \tag{6.4}$$

which implies the coefficient of inequality $\sum_{m=1}^{\infty} |a_m - b_m| \leq \delta \quad (m \in \mathbb{N})$.

Since $g \in \Sigma_p(\alpha, \beta)$, we have $\sum_{m=1}^{\infty} b_m = \frac{1-\alpha}{3+2\beta-\alpha}$.

So that $\left|\frac{f(z)}{g(z)} - 1\right| < \frac{\sum_{m=1}^{\infty} |a_m - b_m|}{1 - \sum_{m=1}^{\infty} b_m} \leq \frac{\delta[3+2\beta-\alpha]}{2(1+\beta)} = 1 - \gamma$, provided γ is given by (6.3).

Hence, by Definition 6.1 , $f \in \Sigma_p(\alpha, \beta, \gamma)$ for γ given by (6.3), which completes the proof of theorem. Q.E.D.

Acknowledgment

The authors express their sincere thanks to the esteemed referee(s) for their careful readings, valuable suggestions and comments, which helped them to improve the presentation of the paper.

References

[1] O. P. Ahuja, G. Murugusundaramoorthy and N. Magesh, *Integral means for uniformly convex and starlike functions associated with generalized hypergeometric functions*, J. Inequal. Pure Appl. Math., 8(4) (2008), 118 -127.

[2] R. Bharathi, R. Parvatham and A. Swaminathan, *On subclasses of uniformly convex functions and corresponding class of starlike functions*, Tamkang J. Maths., 28(1) (1997), 17-32.

[3] J. Clunie, *On meromorphic schlicht functions*, J. London Math. Soc., 34 (1959), 215-216.

[4] A. W. Goodman, *Univalent functions and non-analytic curves*, Proc. Amer. Math. Soc., (1957), 598-601.

[5] A. W. Goodman, *On uniformly convex functions*, Ann. Pol. Math., 56 (1991), 87-92.

[6] A. W. Goodman, *On Uniformly starlike functions*, J. of Math. Anal. and Appl., 155 (1991), 364-370 .

[7] O. P. Juneja and T. R. Reddy, *Meromorphic starlike univalent functions with positive coefficients*, Ann. Univ. Maiiae Curie Sklodowska - Sect. A, 39 (1985), 65-76.

[8] S. Kanas and A. Wisniowska, *Conic regions and k-uniform convexity*, Comput. Appl. Math., 105 (1999), 327-336.

[9] S. Kanas and A. Wisniowska, *Conic domains and starlike functions*, Rev. Roum. Math. Pures Appl., 45 (2000), 647-657.

[10] N. Magesh, N.B.Gatti and S.Mayilvaganan, *On certain class of meromorphic functions with positive and fixed second coefficients involving Liu-Srivastava linear operator*, ISRN Math. Anal., 2012 (2012), Article ID 698307, 11 pages.

[11] S. Mayilvaganan, N. Magesh and N. B. Gatti, *On Certain Subclasses of Meromorphic Functions with Positive Coefficients associated with Liu-Srivastava Linear Operator*, Int. J. of Pure and Applied Math., 113(13) (2017), 132 –141.

[12] G. Murugusundaramoorthy and N. Magesh, *Certain subclasses of starlike functions of complex order involving generalised hypergeometric functions*, Int. J. Math. Math. Sci., Article ID 178605 (2010), 12 pages.

[13] Ch. Pommerenke, *On meromorphic starlike functions*, Pacfic J. Math., 13 (1963), 221-235.

[14] F.Rønning, *On starlike functions associated with parabolic regions*, Ann. Univ. Mariae Curie-Sklodowska- Sect. A, 45 (1991), 117-122.

[15] F. Rønning, *Uniformly convex functions and a corresponing class of starlike functions*, Proc. Amer. Math. Soc., 118 (1993), 189-196.

[16] F. Rønning, *Integral represention of bounded starlike functions*, Ann. Pol. Math, 60(3) (1995), 298-297.

[17] W. C. Royster, *Meromorphic starlike multivalent functions*, Trans. Amer. Math. Soc., 107 (1963), 300-308.

[18] St. Ruscheweyh, *Neighbourhoods of univalent functions*, Proc. Amer. Math. Soc., 81 (1981), 521-527.

[19] H. Silverman, *Univalent functions with negative coefficients*, Proc. Amer. Math. Soc., 51 (1975), 109-116.

[20] B. Venkateswarlu, P. Thirupathi Reddy and N. Rani, *Certain subclass of meromorphically uniformly convex functions with positive coefficients*, Mathematica (Cluj), 61 (84) (1) (2019), 85–97 .